

Two dimensional periodic box-ball system and its fundamental cycle

Shinsuke Iwao

Graduate School of Science, Rikkyo University,
3-34-1 Nishi-Ikebukuro, Toshima-ku, Tokyo, JAPAN

January 12, 2013

Abstract

We study a 2-dimensional Box-Ball system which is a ultradiscrete analog of the discrete KP equation. We construct an algorithm to calculate the fundamental cycle, which is an important conserved quantity of the 2-dim. Box-Ball system with periodic boundary condition, by using the tropical curve theory.

1 Introduction

Ultradiscretization is a limiting procedure by which one can obtain piecewise linear equations from algebraic equations. This procedure allows us to make various piecewise linear dynamical systems, that are called *ultradiscrete integrable systems*, from known discrete integrable systems. The first example of ultradiscrete integrable systems is the Takahashi-Satsuma Box-Ball system (BBS), discovered in 1990 [TS], which is a dynamical system of balls in a one dimensional array of boxes. The BBS shows both a feature of cellular automata and that of solitons.

After the discovery of the BBS, many researchers discovered various kinds of variants of the BBS, such as the BBS with carrier [TM], the BBS with colored balls [TNS] and several kinds of 2-dimensional BBS (2dBBS) [HIK, IH, MNSTTTM].

Around the same time, another relation between the BBS and the solvable quantum model was recognized [HHIKTT]. Both the ultradiscretization and the solvable quantum models are regarded as origins of ultradiscrete integrable systems.

In this paper, we study a 2-dimensional BBS (2dBBS) which is a ultradiscretization of the *discrete KP equation* (dKP)

$$-\delta_m f_{n,m}^{t+1} f_{n+1,m+1}^t + (1 + \delta_m) f_{n+1,m}^t f_{n,m+1}^{t+1} - f_{n,m+1}^t f_{n+1,m}^{t+1} = 0, \quad (1)$$

with a certain periodic boundary condition. Introducing new variables

$$I_{n,m}^t := (1 + \delta_m) \frac{f_{n,m}^t f_{n,m+1}^{t+1}}{f_{n,m+1}^t f_{n,m}^{t+1}}, \quad V_{n,m}^t := \delta_m \frac{f_{n,m}^t f_{n+1,m+1}^t}{f_{n+1,m}^t f_{n,m+1}^t},$$

we can rewrite dKP as $\begin{cases} I_{n,m+1}^t + V_{n,m}^{t+1} = I_{n+1,m}^t + V_{n,m+1}^t \\ I_{n,m}^t V_{n,m}^{t+1} = I_{n+1,m}^t V_{n,m}^t \end{cases}$. Further, we can transform this system as follows:

Lemma 1.1 *The following system implies dKP:*

$$I_{n,m}^t = V_{n,m}^t \left\{ 1 + \left[\frac{V_{n,m}^t}{I_{n+1,m-1}^t} + \frac{V_{n,m}^t V_{n,m-1}^t}{I_{n+1,m-1}^t I_{n+1,m-2}^t} + \frac{V_{n,m}^t V_{n,m-1}^t V_{n,m-2}^t}{I_{n+1,m-1}^t I_{n+1,m-2}^t I_{n+1,m-3}^t} + \dots \right]^{-1} \right\} \quad (2)$$

$$I_{n,m}^t V_{n,m}^{t+1} = I_{n+1,m}^t V_{n,m}^t. \quad (3)$$

Proof. Let $D_m := \frac{V_{n,m}^t}{I_{n+1,m-1}^t} + \frac{V_{n,m}^t V_{n,m-1}^t}{I_{n+1,m-1}^t I_{n+1,m-2}^t} + \frac{V_{n,m}^t V_{n,m-1}^t V_{n,m-2}^t}{I_{n+1,m-1}^t I_{n+1,m-2}^t I_{n+1,m-3}^t} + \dots$. Then D_m satisfies $D_m = \frac{V_{n,m}^t}{I_{n+1,m-1}^t} (1 + D_{m-1})$. This equation and (2-3) imply $I_{n,m+1}^t = V_{n,m+1}^t (1 + D_{m+1}^{-1}) = V_{n,m+1}^t + I_{n+1,m}^t - V_{n,m}^{t+1}$. ■

Under the periodic boundary condition $I_{n,m}^t \equiv I_{n+N,m}^t \equiv I_{n,m+M}^t$, $V_{n,m}^t \equiv V_{n+N,m}^t \equiv V_{n,m+M}^t$, the quantities $\alpha = \prod_{m=1}^M I_{n,m}^t = \prod_{m=1}^M (1 + \delta_m)$, $\beta = \prod_{m=1}^M V_{n,m}^t = \prod_{m=1}^M \delta_m$ should be invariant under the shifts $n \mapsto n+1$, $t \mapsto t+1$. Assume $\alpha > \beta > 0$. Then, (2) is equivalent to

$$I_{n,m}^t = V_{n,m}^t \left(1 + \frac{1 - \beta/\alpha}{\frac{V_{n,m}^t}{I_{n+1,m-1}^t} + \frac{V_{n,m}^t V_{n,m-1}^t}{I_{n+1,m-1}^t I_{n+1,m-2}^t} + \dots + \frac{V_{n,m}^t V_{n,m-1}^t \dots V_{n,m+1}^t}{I_{n+1,m-1}^t I_{n+1,m-2}^t \dots I_{n+1,m}^t}} \right). \quad (4)$$

If all $I_{n,m}^t$, $V_{n,m}^t$ are positive, we introduce a new positive parameter $\varepsilon > 0$ and the variable transformation: $I_{n,m}^t = e^{-\frac{Q_{n,m}^t}{\varepsilon}}$, $V_{n,m}^t = e^{-\frac{W_{n,m}^t}{\varepsilon}}$, $\alpha = k_1 e^{-\frac{A}{\varepsilon}}$, $\beta = k_2 e^{-\frac{B}{\varepsilon}}$. The 2dBBS is the following piecewise linear system which is obtained by taking the limit $\varepsilon \rightarrow 0^+$ of the equations (3, 4):

$$\begin{cases} Q_{n,m}^t = W_{n,m}^t + \min[0, X_{n,m}^t], & W_{n,m}^{t+1} = Q_{n+1,m}^t + W_{n,m}^t - Q_{n,m}^t \\ X_{n,m}^t := \max_{k=0}^{M-1} [\sum_{l=0}^k (Q_{n+1,m-l-1}^t - W_{n,m-l}^t)] \end{cases}. \quad (5)$$

We note $A = \sum_{m=1}^M Q_{n,m}^t$, $B = \sum_{m=1}^M W_{n,m}^t$.

1.1 Examples of 2dBBS

We can regard 2dBBS as a dynamical system $\{W_{n,m}^t\}_{n,m} \mapsto \{W_{n,m}^{t+1}\}_{n,m}$. (See figure 1). Although the




Figure 1: The time evolution rule of 2dBBS.

existence of $\{Q_{n,m}^t\}_{n,m}$ satisfying the periodic boundary condition $Q_{n+M,m}^t = Q_{n,m}^t$ is non-trivial, we can prove the following statement:

Proposition 1.2 *For any $\{W_{n,m}^t\}_{n,m} \in \mathbb{R}^{N \times M}$ and real number A , there uniquely exists $\{Q_{n,m}^t\}_{n,m}$ satisfying the periodic boundary condition and $A = \sum_{m=1}^M Q_{n,m}^t \forall n$.*

Proof. We give the proof in the next section (Remark 2.1). ■

We express the state of the 2dBBS by laying out the following data on two dimensional space: (i) $N \times M$ numbers $\{W_{n,m}^t\}_{n,m}$, (ii) M numbers $\{Q_{1,m}^t\}_m$ as figure 2. This expression is similar to the graphical interpretation of the 2dBBS which is introduced in [HIK, IH].

The following example shows the collision of two solitons ($A = 1, B = 3$).

$Q_{1,M}^t$	$W_{N,M}^t$	$\cdots$	$W_{3,M}^t$	$W_{2,M}^t$	$W_{1,M}^t$	M
$\vdots$	$\vdots$		$\vdots$	$\vdots$	$\vdots$	$\vdots$
$Q_{1,2}^t$	$W_{N,2}^t$	$\cdots$	$W_{3,2}^t$	$W_{2,2}^t$	$W_{1,2}^t$	2
$Q_{1,1}^t$	$W_{N,1}^t$	$\cdots$	$W_{3,1}^t$	$W_{2,1}^t$	$W_{1,1}^t$	$1 = m$
	$n = N$	$\cdots$	3	2	1	

Figure 2: The state of 2dBBS at time t .

. . . . 1.2. . .	 111. .	 2.1.	 11.1
. . . . 1.2. . .	 12. . . .	 3. . . .	 21. . .
. . . 1.2. . . .	. . . 12.	 3.	 21. . . .
1 322.11333	1 332. .2233	1 333. .1323	1 3331.1232
$t=0$	$t=1$	$t=2$	$t=3$

This paper is arranged as follows: In §2, we review the connection between the dKP and the algebraic geometry. We introduce the *spectral curve* of the dKP, that is an important invariant of the equation. In §3, we review several basic results of the tropical geometry according to [I1, IMS, M, MZ]. In §4, we obtain a tropical analog of the spectral curve of 2dBBS and a tropical-geometric expression of the invariants. Moreover, we have an algorithm to calculate the fundamental cycle of 2dBBS.

Notations: For a positive integer M and a ring R with 1, $\text{Mat}(M, R)$ denotes the R -algebra of matrices over R of size M . E denotes the identity matrix and $E_{i,j}$ denotes the matrix of which (i, j) -element is 1, and other elements are 0. We denote by $\text{diag}(a_1, \dots, a_M)$ the diagonal matrix of which (i, i) -element equals to a_i . Define $S := (\delta_{i+1,j})_{i,j} + y \cdot E_{M,1} \in \text{Mat}(M, \mathbb{C}[y])$.

Let $\mathcal{O}$ be a $\mathbb{C}[y^{\pm 1}]$ -algebra and $\mathcal{O}^\infty[M]$ be the $\mathcal{O}$ -module defined by

$$\mathcal{O}^\infty[M] := \{(\cdots, a_{-1}, a_0, a_1, \dots) \mid a_i \in \mathcal{O}, a_{i+M} = y \cdot a_i\},$$

which is isomorphic to $\mathcal{O}^M = \mathcal{O} \times \cdots \times \mathcal{O}$ (M times) as $\mathcal{O}$ -modules through the isomorphism $\mathcal{O}^\infty[M] \rightarrow \mathcal{O}^M; (a_i)_{i \in \mathbb{Z}} \mapsto (a_i)_{i=1}^M$. An element of $\mathcal{O}^\infty[M]$ is called a M -periodic vector.

The set of $\mathcal{O}$ -automorphisms of $\mathcal{O}^M$ is naturally identified as $\text{Mat}(M, \mathcal{O})$. Similarly, the set of $\mathcal{O}$ -automorphisms of $\mathcal{O}^\infty[M]$ is identified as the $\mathcal{O}$ -submodule P of $\text{Mat}(\infty, \mathbb{C})$ which is the image of the morphism

$$\text{Mat}(M, \mathbb{C}[y^{\pm 1}]) \rightarrow \text{Mat}(\infty, \mathbb{C}); (z_{i,j})_{i,j=1}^M \mapsto (\tilde{z}_{i,j})_{i,j \in \mathbb{Z}},$$

where $z_{i,j} = \sum_{n \in \mathbb{Z}} \tilde{z}_{i,j+nM} \cdot y^n$ ($1 \leq i, j \leq M$), $\tilde{z}_{i+M,j} = \tilde{z}_{i,j}$. An element of P is called a M -periodic matrix. For a matrix $Z = (z_{i,j})_{i,j=1}^M \in \text{Mat}(M, \mathcal{O})$, we express the associated M -periodic matrix as $\tilde{Z} = (\tilde{z}_{i,j})_{i,j \in \mathbb{Z}}$. Similarly, for $a = (a_i)_{i=1}^M \in \mathcal{O}^M$, we denote the associated M -periodic vector by $\tilde{a} = (\tilde{a}_i)_{i \in \mathbb{Z}}$. It follows that $(\tilde{Z} \cdot a) = \tilde{Z} \cdot \tilde{a}$.

2 Periodic discrete KP equation

2.1 spectral curve

Periodic discrete KP equation (Lemma 1.1) is rewritten as the following matrix form:

$$L_n^{t+1}(y)R_n^t(y) = R_{n+1}^t(y)L_n^t(y), \quad (6)$$

where $L_n^t(y) = \text{diag}(V_{n,1}^t, \dots, V_{n,M}^t) + S$, $R_n^t(y) = \text{diag}(I_{n,1}^t, \dots, I_{n,M}^t) + S$. By using the new matrix

$$X_n^t(y) := L_{n+N-1}^t \cdots L_{n+1}^t L_n^t, \quad (L_n^t = L_n^t(y)), \quad (7)$$

we can transform the dKP to the following matrix form:

$$X_n^{t+1} R_n^t = R_n^t X_n^t, \quad \text{or equivalently,} \quad X_{n+1}^t L_n^t = L_n^t X_n^t. \quad (8)$$

Form (8), the characteristic polynomial $\Phi(x, y) := \det(X_n^t(y) - xE)$ should be invariant under $t \mapsto t+1$, $n \mapsto n+1$. Let $\tilde{C} \subset \mathbb{P}^1 \times \mathbb{P}^1$ be the algebraic curve defined by Φ and $C_0 \subset \mathbb{C}^2$ be the affine part of $\tilde{C}$. Put $\eta = x^{-1}$, $\zeta = y^{-1}$, $d := \text{g.c.d.}(N, M)$, $N = dN_1$ and $M = dM_1$. By direct calculations, we have the following expression [MIT]:

$$\eta^M \zeta^N \cdot \Phi = (\eta^{M_1} - \zeta^{N_1})^d + \sum_{NM < Ni + Mj \leq 2NM, i, j \geq 0} c_{i,j} \eta^i \zeta^j, \quad c_{i,j} \neq 0, \quad (9)$$

which implies that the set $\tilde{C} \setminus C_0$ consists of only one point $p_0 : (\eta, \zeta) = (0, 0)$.

By (9), we can introduce a coordinate variable k such that $\eta = k^N \cdot (1 + c_1 k + c_2 k^2 + \dots)$ and $\zeta = k^M$, where $c_1 \neq 0$. Denote by C the analytic curve obtained from $\tilde{C}$ of which the local coordinate at p_0 is k . We call C a *spectral curve* of the dKP.

Let us define the following two sets:

$$\begin{aligned} \mathcal{V}(\beta) &:= \{(V_{n,m})_{n,m} \in \mathbb{C}^{N \times M} \mid \beta = \prod_m V_{n,m} \ (\forall n)\}, \\ \mathcal{W}(\beta) &:= \{X \mid X = L_N \cdots L_2 L_1, \ L_n = \text{diag}(V_{n,1}, \dots, V_{n,M}) + S, \ (V_{n,m}) \in \mathcal{V}(\beta)\}. \end{aligned}$$

Let $j : \mathcal{V}(\beta) \rightarrow \mathcal{W}(\beta)$ be the natural surjection which sends an element $(V_{n,m})_{n,m} \in \mathcal{V}(\beta)$ to the matrix $X = L_N \cdots L_2 L_1$, $L_n = \text{diag}(V_{n,1}, \dots, V_{n,M}) + S$. One can prove that j is bijective (Lemma 3.1. [KNY]). We will often identify $\mathcal{V}(\beta)$ with $\mathcal{W}(\beta)$.

Through this procedure, we have the following map:

$$\text{SC} : \mathcal{V}(\beta) \cong \mathcal{W}(\beta) \rightarrow \{\text{spectral curves of dKP}\}. \quad (10)$$

The *isolevel set associated with a spectral curve C* is the inverse image $\mathcal{T}_C := \text{SC}^{-1}(C)$.

For two matrices $X, Y \in \mathcal{T}_C$, we say $X \sim Y$ if there exists an invertible diagonal matrix D such that $X = DYD^{-1}$. This defines an equivalence relation over $\mathcal{T}_C$. We write the quotient set induced by this relation as $\mathcal{R}_C := \mathcal{T}_C / \sim$.

2.2 time evolution of dKP

Unfortunately, the equation $X_n^{t+1} R_n^t = R_n^t X_n^t$ (8) does not define the time evolution $X_n^t \mapsto X_n^{t+1}$ uniquely. In fact, the following proposition can be proved.

Proposition 2.1 *For given $X_n^t \in \mathcal{W}(\beta)$, there exist M pairs $(R^{(1)}, X^{(1)}), \dots, (R^{(M)}, X^{(M)})$ such that (i) $R^{(i)} = \text{diag}(I_1^{(i)}, \dots, I_M^{(i)}) + S$, (ii) $\alpha = \prod_{m=1}^M I_m^{(i)}$, (iii) $X^{(i)} \in \mathcal{W}(\beta)$, (iv) $X^{(i)} R^{(i)} = R^{(i)} X_n^t$.*

However, we can uniquely specify a certain time evolution with an additional condition.

Proposition 2.2 *Let $X_n^t = L_{n+N-1} \cdots L_{n+1} L_n \in \mathcal{W}(\beta)$, $L_n^t = \text{diag}(V_{n,1}^t, \dots, V_{n,M}^t) + S$ such that $V_{n,m}^t > 0$. Then, there uniquely exist two matrices X_n^{t+1} , R_n^t such that (i) $R_n^t = \text{diag}(I_1^t, \dots, I_M^t) + S$, (ii) $\alpha = \prod_{m=1}^M I_m^t$, (iii) $X_n^{t+1} \in \mathcal{W}(\beta)$, (iv) $X_n^{t+1} R_n^t = R_n^t X_n^t$, (v) $I_{n,m}^t > 0$.*

By this proposition, we can say that a certain type of the time evolution of X_n^t is determined by (8). We will give the proofs of propositions 2.1, 2.2 in the appendix.

Remark 2.1 *Proposition 2.1 implies proposition 1.2.*

2.3 linearisation

The method to solve the initial value problem of discrete integrable systems with a spectral curve have been established by many authors. We review some fundamental results concerning the *linearisation*.

We consider the three automorphisms T_t, T_n, T_m of $\mathcal{T}_C$ defined by

$$T_t(X_n^t) := X_n^{t+1} = R_n^t X_n^t (R_n^t)^{-1}, \quad T_n(X_n^t) := X_{n+1}^t = L_n^t X_n^t (L_n^t)^{-1}, \quad (11)$$

$$T_m(X_n^t) := X_n^t|_{m \rightarrow m+1} = S X_n^t S^{-1}, \quad (12)$$

where T_t is the unique time evolution defined in §2.2. Let $X_{n,m}^t := S^{m-1} X_n^t S^{-m+1}$. Then it follows that $X_{n,m+1}^t = T_m(X_{n,m}^t)$.

For an algebraic curve C , we write $\text{Div } C := \bigoplus_{p \in C} \mathbb{Z} \cdot p$. The Picard group $\text{Pic } C$ is the quotient group of $\text{Div } C$ induced by the linear equivalence relation. Denote by $\text{Pic}^n C$ the subset of $\text{Pic } C$ consisted of the divisors of degree n . For a rational function f over C , $(f)_0$ (*resp.* $(f)_\infty$) denotes the divisor of zeros (*resp.* poles) of f . The following theorem 2.3 is a fundamental result concerning the algebro-geometric aspects of spectral curves proved in [MM].

Theorem 2.3 *There exists a mapping $\tilde{\varphi} : \mathcal{R}_C \rightarrow \text{Div}^g C$ such that:*

- (i) $\tilde{\varphi}(X)$ is a general divisor of degree g for any $X \in \mathcal{R}_C$,
- (ii) the induced mapping $\varphi : \mathcal{R}_C \rightarrow \text{Pic}^g C$ is injective,
- (iii) we have $\tilde{\varphi}(X) = (f_1/f_M)_0 - (M-1)p_0$, where $f_i = f_i(x, y)$ is the i -th component of a eigenvector of $X = X(y)$ belonging to an eigenvalue x . ■

Let $X_n^t \in \mathcal{T}_C$. We denote by $\mathbf{v}_n^t = (f_{n,1}^t, \dots, f_{n,M}^t)^T$ a eigenvector of X_n^t belonging to x i.e.

$$X_n^t \mathbf{v}_n^t = x \mathbf{v}_n^t. \quad (13)$$

By (8), we have the following evolution equation

$$\mathbf{v}_n^{t+1} = R_n^t \mathbf{v}_n^t, \quad \mathbf{v}_{n+1}^t = L_n^t \mathbf{v}_n^t. \quad (14)$$

Equations (13, 14) can be regarded as linear equations over the rational function field $\mathcal{K}$ over C . Naturally, $\mathcal{K}$ has a structure of $\mathbb{C}[y^{\pm 1}]$ -algebra. Using M -periodic matrices and vectors, we rewrite these equations as

$$\tilde{X}_n^t \tilde{\mathbf{v}}_n^t = x \tilde{\mathbf{v}}_n^t, \quad \tilde{\mathbf{v}}_n^{t+1} = \tilde{R}_n^t \tilde{\mathbf{v}}_n^t, \quad \tilde{\mathbf{v}}_{n+1}^t = \tilde{L}_n^t \tilde{\mathbf{v}}_n^t. \quad (15)$$

By definition of X_n^t , the M -periodic matrix $\tilde{X}_n^t$ is expressed as $\tilde{X}_n^t = \tilde{S}^N + \tilde{Z}$, where $\tilde{Z} = (\tilde{z}_{i,j})_{i,j}$ is a M -periodic matrix such that $\tilde{z}_{i,j} \neq 0 \Rightarrow 0 \leq j - i < N$. Then, the first equation of (15) is rewritten as $\tilde{S}^N \tilde{\mathbf{v}}_n^t = (x \tilde{E} - \tilde{Z}) \tilde{\mathbf{v}}_n^t$. Equivalently, we have

$$\tilde{f}_{n,m+N}^t = x \tilde{f}_{n,m}^t - \sum_{k=0}^{N-1} \tilde{z}_{m,m+k} \tilde{f}_{n,m+k}^t. \quad (16)$$

Let $\mathbf{w}_n^t = (\tilde{f}_{n,1}^t, \dots, \tilde{f}_{n,N}^t)^T$, $M_n^t = \text{diag}(I_{n,1}^t, \dots, I_{n,N}^t) + U_1$ and $H_n^t = \text{diag}(V_{n,1}^t, \dots, V_{n,N}^t) + U_1$, where $U_m = (\delta_{i+1,j})_{i,j} + (x - \tilde{z}_{m,m}) E_{N,1} - \tilde{z}_{m,m+1} \cdot E_{N,2} - \dots - \tilde{z}_{m,m+N-1} \cdot E_{N,N}$. Then, from (16), we can rewrite (14) as

$$\mathbf{w}_n^{t+1} = M_n^t \mathbf{w}_n^t, \quad \mathbf{w}_{n+1}^t = H_n^t \mathbf{w}_n^t. \quad (17)$$

Theorem 2.4 ([I2]) (i) *There exist divisors $\mathcal{T}, \mathcal{N}, \mathcal{M}_1, \mathcal{M}_2, \dots, \mathcal{M}_M \in \text{Pic}(C)$ of degree 0 such that $\varphi(X_{n,m}^{t+1}) = \varphi(X_{n,m}^t) + \mathcal{T}$, $\varphi(X_{n+1,m}^t) = \varphi(X_{n,m}^t) + \mathcal{N}$, $\varphi(X_{n,m+1}^t) = \varphi(X_{n,m}^t) + \mathcal{M}_m$.*

(ii) *The divisors $\mathcal{T}, \mathcal{N}, \mathcal{M}_m$ are expressed as follows:*

$$\mathcal{T} = p_{\mathcal{T}}^+ - p_{\mathcal{T}}^-, \quad \mathcal{N} = p_{\mathcal{N}}^+ - p_{\mathcal{N}}^-, \quad \mathcal{M}_m = p_{\mathcal{M}}^+(m) - p_{\mathcal{M}}^-(m), \quad (18)$$

where $p_{\mathcal{T}}^{\pm}, p_{\mathcal{N}}^{\pm}, p_{\mathcal{M}}^{\pm}(m)$ are certain points on C .

(iii) The x, y coordinates of $p_{\mathcal{T}}^{\pm}, p_{\mathcal{N}}^{\pm}, p_{\mathcal{M}}^{\pm}(m)$ are expressed as

$$\begin{aligned} x_{\mathcal{T}}^+ &= (\det M_n^t)_{\infty}, & x_{\mathcal{T}}^- &= (\det M_n^t)_0, & y_{\mathcal{T}}^+ &= (\det R_n^t)_{\infty}, & y_{\mathcal{T}}^- &= (\det R_n^t)_0, \\ x_{\mathcal{N}}^+ &= (\det H_n^t)_{\infty}, & x_{\mathcal{N}}^- &= (\det H_n^t)_0, & y_{\mathcal{N}}^+ &= (\det L_n^t)_{\infty}, & y_{\mathcal{N}}^- &= (\det L_n^t)_0, \\ x_{\mathcal{M}}^+(m) &= (\det U_m)_{\infty}, & x_{\mathcal{M}}^-(m) &= (\det U_m)_0, \\ y_{\mathcal{M}}^+(m) &= (\det S)_{\infty}, & y_{\mathcal{M}}^-(m) &= (\det S)_0. \quad \blacksquare \end{aligned}$$

Roughly speaking, theorem 2.4 states that the determinants of matrices $L_n^t, R_n^t, H_n^t, M_n^t, \dots$ characterize the action of T_n, T_m, T_t .

Lemma 2.5 $\det R_n^t = \alpha - (-1)^M y$, $\det L_n^t = \beta - (-1)^M y$, $\det H_n^t = (-1)^{N+1} x$, $\det S = (-1)^{M+1} y$, $\det U_m = (-1)^{N+1} (x - V_{N,m}^t \cdots V_{2,m}^t V_{1,m}^t)$, $\det M_n^t = (-1)^N (x - \kappa)$, where κ is a minimum eigenvalue of $X_n^t|_{y=(-1)^M \alpha}$ in absolute value.

Proof. We prove these lemmas in the appendix. $\blacksquare$

From theorem 2.4 and lemma 2.5, the representations of the actions T_t, T_n, T_m on $\text{Pic}^g C$ are completely determined:

Proposition 2.6 The divisors $\mathcal{T}, \mathcal{N}, \mathcal{M}_m$ in theorem 2.4 are expressed as $\mathcal{T} = p_0 - p_1$, $\mathcal{N} = p_0 - p_2$, $\mathcal{M}_m = p_0 - p_3^{(m)}$, where $p_1 : (x, y) = (\kappa, \alpha)$, $p_2 : (x, y) = (0, \beta)$, $p_3^{(m)} : (x, y) = (\gamma_m, 0)$, $\gamma_m = V_{N,m}^t \cdots V_{2,m}^t V_{1,m}^t$.

2.4 $\mathcal{T}_C$ and $\mathcal{R}_C$

Theorem 2.4 states that the evolutions $t \mapsto t+1, n \mapsto n+1, m \mapsto m+1$ over $\mathcal{R}_C$ can be linearized on $\text{Pic}^g(C)$ through the injection $\varphi : \mathcal{R}_C \rightarrow \text{Pic}^g(C)$. To lift this result to the isolevel set $\mathcal{T}_C$, we will study the structures of the two sets $\mathcal{R}_C$ and $\mathcal{T}_C$. Let $d := \text{g.c.d.}(N, M)$.

Our aim in this section is to construct a bijection

$$\mathcal{T}_C \rightarrow \mathcal{R}_C \times (\mathbb{C}^{\times})^{d-1} \quad (19)$$

which has some nice behavior. Denote by $[X]$ the image of $X \in \mathcal{T}_C$ through the natural projection $\mathcal{T}_C \rightarrow \mathcal{R}_C = \mathcal{T}_C / \sim$. If $[X] = [Y]$, there exists a diagonal invertible matrix D such that $X = DYD^{-1}$.

Lemma 2.7 Let D be a diagonal and invertible matrix. Then, $X, DXD^{-1} \in \mathcal{T}_C \iff S^d DS^{-d} = D$.

Proof. $\Rightarrow$) Because any element $X \in \mathcal{T}_C$ is written as $X = S^N + \sum_{i=0}^{N-1} X_i S^i$, (X_i are diagonal) uniquely, we have $S^N DS^{-N} = D$. On the other hand, by definition of the matrix S , it follows that $S^M DS^{-M} = D$. Due to these equations, we have $S^d DS^{-d} = D$. $\Leftarrow$) Any element $X \in \mathcal{T}_C$ is written as $X = (L_N + S) \cdots (L_2 + S)(L_1 + S)$. Putting $L'_n := (S^{-n} DS^n) L_n (S^{n-1} D^{-1} S^{-n+1})$ and $X' := (L'_N + S) \cdots (L'_1 + S)$, we have $X' = DXD^{-1} \in \mathcal{T}_C$. $\blacksquare$

Let $G(\cong (\mathbb{C}^{\times})^d)$ be the multiplicative group of diagonal matrices defined by

$$G := \{D \mid D : \text{invertible, diagonal, } S^d DS^{-d} = D\},$$

and $H(\cong \mathbb{C}^{\times}) \subset G$ be the normal subgroup $H := \{cE \mid c \in \mathbb{C}^{\times}\}$. By lemma 2.7, the quotient group G/H acts on $\mathcal{T}_C$ by $G/H \ni D \mapsto \{X \mapsto DXD^{-1}\} \in \text{End}(\mathcal{T}_C)$. Because the group action is free, the orbit $(G/H) \cdot X$ is isomorphic to $G/H \cong (\mathbb{C}^{\times})^{d-1}$ for any $X \in \mathcal{T}_C$.

Therefore, any map $i : \mathcal{T}_C \rightarrow G/H$ satisfying $i(DXD^{-1}) = D \cdot i(X)$ induces a bijection $\mathcal{T}_C \rightarrow \mathcal{R}_C \times (G/H)$. To define a good bijection, we focus on an eigenvector of X at the point $p_0 \in C$. Let k be the local coordinate around $p_0 \in C$.

Lemma 2.8 *Let X be an element of $\mathcal{T}_C$, $\mathcal{K}$ be the rational function field of C . Assume $\mathbf{v} \in \mathcal{K}^M$ satisfies $X\mathbf{v} = x\mathbf{v}$. Up to multiplication by a constant, the i -th component f_i of $\mathbf{v}$ has the following expression near p_0 :*

$$f_i = a_i k^{M-i} + o(k^{M-i}), \quad a_i \neq 0.$$

Moreover, we have $a_{i+d} = a_i$ for all i .

Proof. See the appendix. ■

Define the map $i : \mathcal{T}_C \rightarrow G/H (\cong (\mathbb{C}^\times)^d / \mathbb{C}^\times)$ by $X \mapsto [a_1 : a_2 : \dots : a_d]^T$, where a_i is the non-zero number in lemma 2.8. By definition of i , we can check $i(DXD^{-1}) = D \cdot i(X)$ immediately. As stated above, this map induces a bijection $\mathcal{T}_C \rightarrow \mathcal{R}_C \times G/H$.

Proposition 2.9 *Let $X_{n,m}^t \in \mathcal{T}_C$ and $i(X_{n,m}^t) = [a_1 : a_2 : \dots : a_d]^T$. Then it follows that $i(X_{n+1,m}^t) = i(X_{n,m+1}^t) = i(X_{n,m}^{t+1}) = [a_2 : a_3 : \dots : a_d : a_1]^T$.*

Proof. Let f_i be the i -th element of $\mathbf{v}$ such that $X_{n,m}^t \mathbf{v} = x\mathbf{v}$, and f'_i be the i -th element of $\mathbf{v}'$ such that $X_{n+1,m}^t \mathbf{v}' = x\mathbf{v}'$. Because $X_{n+1,m}^t = (S^{m-1}L_n^t S^{-m+1})X_{n,m}^t(S^{m-1}L_n^t S^{-m+1})^{-1}$, we have $\mathbf{v}' = (S^{m-1}L_n^t S^{-m+1})\mathbf{v}$, which implies $f'_i = \begin{cases} f_{i+1} + V_{n,m+i}^t f_i & i \neq M \\ y f_1 + V_{n,m}^t f_M & i = M \end{cases}$. From this equation and lemma 2.8, we have $f'_i = \begin{cases} a_{i+1}k^{M-i-1} + o(k^{M-i-1}) & i \neq M \\ a_1 k^{-1} + o(k^{-1}) & i = M \end{cases}$. By multiplying k to all f'_i simultaneously, we obtain the desired expression. The equations about $X_{n,m+1}^t$ and $X_{n,m}^{t+1}$ are proved similarly. ■

Let φ be the injection $\mathcal{R}_C \rightarrow \text{Pic}^g(C)$ introduced in §2.3, and $\mathbf{a} := \text{Im}\varphi$ be the image of φ . From theorem 2.4 and proposition 2.9, we have :

Theorem 2.10 *There exist divisors $\mathcal{T}, \mathcal{M}_m, \mathcal{N} \in \text{Pic}^0(C)$ and a bijection $\eta : \mathcal{T}_C \rightarrow \mathbf{a} \times G/H$ such that: For $\eta(X_{n,m}^t) = (\mathcal{D}, \mathbf{a})$,*

$$\begin{aligned} \eta(X_{n+1,m}^t) &= (\mathcal{D} + \mathcal{T}, \sigma(\mathbf{a})), & \eta(X_{n,m+1}^t) &= (\mathcal{D} + \mathcal{M}_m, \sigma(\mathbf{a})), \\ \eta(X_{n,m}^{t+1}) &= (\mathcal{D} + \mathcal{N}, \sigma(\mathbf{a})), & \sigma([a_1 : a_2 : \dots : a_d]^T) &:= [a_2 : a_3 : \dots : a_d : a_1]^T. \end{aligned}$$

Corollary 2.11 $\mathcal{T}_C$ is embedded into $\text{Jac } C \times (\mathbb{C}^\times)^{d-1}$, where $\text{Jac } C$ is the Jacobi variety of C .

3 Review of tropical geometry

We briefly review basic results of the tropical geometry. For details, see [IMS, MZ].

3.1 ultradiscretization of functions

Let $\mathbb{R}_+^L := (\mathbb{R}_{>0})^L$ be the set of L -positive vectors and $T = (0, \infty)$ be the open interval of infinite length. For a topological set A , denote by $\text{Hom}(T, A)$ the set of continuous maps from T to A . Any continuous map $f : A \rightarrow B$ induces a mapping $f^* : \text{Hom}(T, A) \rightarrow \text{Hom}(T, B)$.

Let A be a subset of $(\mathbb{C}^\times)^L$. We define the subset $\mathcal{U}^A \subset \text{Hom}(T, A)$ by

$$\mathcal{U}^A := \{(s_1, \dots, s_L) \in \text{Hom}(T, A) ; \lim_{\varepsilon \rightarrow 0+} |\varepsilon \log |s_i(\varepsilon)|| < +\infty\}. \quad (20)$$

Let $\text{UD} : \mathcal{U}^A \rightarrow \mathbb{R}^L$ be the map expressed as $(s_i)_i \mapsto (-\lim_{\varepsilon \rightarrow 0+} \varepsilon \log |s_i(\varepsilon)|)_i$.

Definition 3.1 A map $f : A \rightarrow B$ is ultradiscretizable if: (i) $f^*(\mathcal{U}^A) \subset \mathcal{U}^B$, (ii) There exists a map $\text{UD}(f) : \mathbb{R}^L \rightarrow \mathbb{R}^L$ such that the following diagram is commutative:

$$\begin{array}{ccc} \mathcal{U}^A & \xrightarrow{f^*} & \mathcal{U}^B \\ \text{UD} \downarrow & & \downarrow \text{UD} \\ \mathbb{R}^L & \xrightarrow{\text{UD}(f)} & \mathbb{R}^L \end{array} .$$

We call the map $\text{UD}(f)$ the *ultradiscretization* of f . It is not easy to determine whether a given function f is ultradiscretizable. Here, we introduce two simple sufficient conditions.

Example 3.1 (Totally positive polynomial) If f is expressed as a totally positive polynomial, then f is ultradiscretizable.

Example 3.2 (Roots of polynomial) Let $A := (\mathbb{C}^\times)^{N+1}$, $B := \mathbb{C}^\times$ and $\Phi(x) = s_N x^N + \dots + s_1 x + s_0$ with $s_i \in \text{Hom}(T, \mathbb{C}^\times)$. Then there exists a continuous function $x = x(\varepsilon)$ such that $\Phi(x(\varepsilon)) = 0$ for all ε . Define $f_\Phi^*(s_0, s_1, \dots, s_N) := x$. Therefore, $f_\Phi : A \rightarrow B$ is ultradiscretizable. In fact, $\text{UD}(f_\Phi)$ is a piecewise linear function of $\text{UD}(s_0), \dots, \text{UD}(s_N)$.

3.2 tropical curves

3.2.1 definition

Define the subset $\mathcal{U} := \mathcal{U}^{\mathbb{C}^\times}$ of $\text{Hom}(T, \mathbb{C}^\times)$ as (20). For a polynomial $\Phi = \sum_{w=(w_1, w_2) \in \mathbb{Z}^2} c_w x^{w_1} y^{w_2} \in \mathcal{U}[x^{\pm 1}, y^{\pm 1}]$, denote $\text{UD}(X, Y; \Phi) := \min_{w \in \mathbb{Z}^2} [\text{UD}(c_w) + w_1 X + w_2 Y]$. A *plane tropical curve* Γ^0 defined by Φ is a subset of $\mathbb{R}^2$ which is expressed as

$$\Gamma^0 := \{(X_0, Y_0) \mid \text{The function } \text{UD}(X, Y; \Phi) \text{ is not smooth at } (X, Y) = (X_0, Y_0)\}.$$

Example 3.3 Let $e := e^{-1/\varepsilon}$ and $\Phi = y^2 + y(x^3 + x^2 + ex + e^2) + e^6$. Then, we have $\text{UD}(X, Y; \Phi) = \min[2Y, Y + 3X, Y + 2X, Y + X + 1, Y + 2, 6]$. The tropical curve Γ^0 is given as figure 3.




Figure 3: The tropical curve defined by $\Phi = y^2 + y(x^3 + x^2 + ex + e^2) + e^6$.

For $P \in \Gamma^0$, we define $\Phi^P := \sum_{w \in \Lambda_\Phi(P)} c_w x^{w_1} y^{w_2}$, where $\Lambda_\Phi(P) := \{w \in \Lambda_\Phi \mid \text{UD}(X_P, Y_P) = \text{UD}(c_w) + w_1 X_P + w_2 Y_P, P = (X_P, Y_P)\}$. Let $\text{mult}(P) := \sharp\{\text{irreducible components of } \Phi^P \in \mathcal{U}[x^{\pm 1}, y^{\pm 1}]\}$.

Definition 3.2 A tropical curve with multiplicity is a non-oriented graph Γ with surjection $\iota : \Gamma \rightarrow \Gamma^0$ such that $\sharp\{\iota^{-1}(P)\} = \text{mult}(P)$ for all $P \in \Gamma^0$.




Figure 4: The tropical curve with multiplicity defined by Φ .

Example 3.4 Let Φ be the polynomial in example 3.3. Then we have, for example, $\Phi^{(1,4)} = \mathbf{e}yx + \mathbf{e}^2y + \mathbf{e}^6$, $\Phi^{(0,3)} = yx^3 + yx^2 + \mathbf{e}yx = yx(x + \xi_+)(x + \xi_-)$, where $\xi_+ = \mathbf{e} + \mathbf{e}^2 + \dots$, $\xi_- = 1 - \mathbf{e} - \dots$. The tropical curve with multiplicity Γ is given as figure 4.

Note that ι is finite and locally homeomorphic without finite numbers of points. We abbreviate the pull-backed coordinates ι^*X and ι^*Y over Γ as X and Y .

3.2.2 tropical integration

Let $\iota : \Gamma \rightarrow \Gamma^0$ be a tropical curve with multiplicity. Because ι is locally homeomorphic without finite numbers of points, we can induce a metric on Γ by pulling back the metric of Γ^0 defined by the lattice length §5 [M].

Denote by $\mathcal{P}$ the free $\mathbb{Z}$ -module generated by oriented paths on Γ . Let $(\cdot, \cdot) : \mathcal{P} \times \mathcal{P} \rightarrow \mathbb{R}$ be the *tropical bilinear form* [M], which is defined by $(\gamma_1, \gamma_2) := (\pm 1) \cdot \|\gamma_1 \cap \gamma_2\|$. Fix a basis $\beta_1, \dots, \beta_g$ of $H_1(\Gamma; \mathbb{Z})$. Then, we obtain the *period matrix* of Γ , which is a $g \times g$ matrix $B_\Gamma = ((\beta_i, \beta_j))_{i,j=1}^g$. For a fixed point $Q_0 \in \Gamma$, we define the *tropical Abel-Jacobi mapping* $F_{Q_0} : \Gamma \rightarrow \mathbb{R}^g / B_\Gamma \mathbb{Z}^g$ by

$$F_{Q_0}(P) := \left((\gamma_{Q_0 \rightarrow P}, \beta_i) \right)_{i=1}^g \pmod{B_\Gamma \mathbb{Z}^g},$$

where $\gamma_{Q_0 \rightarrow P} \in \mathcal{P}$ is a path from Q_0 to P . We call the variety $J(\Gamma) := \mathbb{R}^g / B_\Gamma \mathbb{Z}^g$ the *tropical Jacobi variety*. The mapping F_{Q_0} can be linearly extended to $\text{Div } \Gamma = \bigoplus_{p \in \Gamma} \mathbb{Z} \cdot p$.

3.3 ultradiscrete limit of Abelian integrals

In this section, we review the theorem in the paper [I1], which explains a certain direct relation between Abelian integrals and tropical integrals. For a polynomial $\Phi(x, y) \in \mathcal{U}[x^{\pm 1}, y^{\pm 1}]$, there exists a holomorphic family of curves $\pi : V \rightarrow T$ such that $\pi^{-1}(\varepsilon)$ is an algebraic curve defined by $\Phi|_\varepsilon \in \mathbb{C}[x^{\pm 1}, y^{\pm 1}]$. Denote by g the genus of these curves.

Let $\alpha_i : A \rightarrow T$ and $\beta_i : B \rightarrow T$ ($i = 1, \dots, g$) be a holomorphic (oriented) subfamily of π such that $\alpha_i^{-1}(\varepsilon)$ and $\beta_i^{-1}(\varepsilon)$ are symplectic basis of $H_1(\pi^{-1}(\varepsilon); \mathbb{Z})$. We call such (α_i, β_i) a *symplectic basis* of π .

The following is the main result of the paper [I1].

Theorem 3.5 ([I1]) *There exists a canonical correspondence*

$$\mathfrak{X} : \{(\alpha_i, \beta_i)_i \mid \text{symplectic basis of } \pi : V \rightarrow T\} \rightarrow \{(\beta_i)_i \mid \text{basis of } H_1(\Gamma; \mathbb{Z})\}$$

and holomorphic 1-forms ω_i ($i = 1, \dots, g$) over V such that

$$\int_{\alpha_i^{-1}(\varepsilon)} \omega_j = \delta_{i,j}, \quad -2\pi i \lim_{\varepsilon \rightarrow 0} \left(\varepsilon \int_{\beta_i^{-1}(\varepsilon)} \omega_j \right) = B_{i,j},$$

where $B_{i,j}$ is the (i,j) -component of B_Γ defined by the basis $\mathfrak{X}(\alpha_i, \beta_i)$. $\blacksquare$

Let $P \in \Gamma$. Because the field $\mathcal{U}$ is algebraically closed, we can find a holomorphic section $p : T \rightarrow V$ of π such that $\text{UD}(x(p(\varepsilon))) = X(P)$, $\text{UD}(y(p(\varepsilon))) = Y(P)$. In this case, we denote ' $p \rightarrow P$ '.

Theorem 3.6 *Let p, q be two sections $T \rightarrow V$. Assume there exist two points $P, Q \in \Gamma$ such that $p \rightarrow P$, $q \rightarrow Q$. Then, the following relations are held*

$$-2\pi i \lim_{\varepsilon \rightarrow 0^+} \varepsilon \cdot f_{q(\varepsilon)}(p(\varepsilon)) = F_Q(P) \in J(\Gamma). \quad \blacksquare$$

4 Application for ultradiscrete systems

From the results in §2 and §3, we obtain the tropical analog of the linearizing theorem for the ultradiscrete integrable systems. Let J be the quotient space $J := (\mathbb{C}^g \times \text{GL}(g, \mathbb{C})) / \sim$, where $(\mathbf{v}, B) \sim (\mathbf{v}', B') \Leftrightarrow B = B'$ and $\mathbf{v} - \mathbf{v}' \in \mathbb{Z}^g + B\mathbb{Z}^g$. The analytical set J is regarded as a set of *all* Jacobi varieties. Any element of J is expressed as $(\mathbf{v} \bmod (\mathbb{Z}^g + B\mathbb{Z}^g), B)$.

4.1 ultradiscretization

4.1.1 family of isolevel sets

For an $\beta \in \text{Hom}(T, \mathbb{C}^\times)$, define

$$\mathcal{V}(\beta) := \left\{ (V_{n,m})_{n=1, m=1}^{N,M} \in \text{Hom}(T, \mathbb{C}^{N \times M}) \mid \begin{array}{l} (i) \beta = \prod_m V_{n,m} \text{ for all } n, \\ (ii) \lim_{\varepsilon \rightarrow 0^+} |\varepsilon \log |V_{n,m}(\varepsilon)|| < +\infty \end{array} \right\}.$$

As in §2.1, any $\{V_{n,m}\}_{n,m} \in \mathcal{V}(\alpha)$ is identified with the matrix function $X_n := L_{n+N-1} \cdots L_{n+1} L_n$, where $L_n = \text{diag}(V_{n,1}, \dots, V_{n,M}) + S$. Let $\pi : V \rightarrow T$ be the holomorphic family of curves defined by $\Phi = \det(X_n - x \cdot \text{id.})$. Consider the correspondence: $\text{SC}^* : \mathcal{V}(\alpha) \rightarrow \{\text{hol. families over } T\}; \{V_{n,m}\}_{n,m} \mapsto \{\pi : V \rightarrow T\}$. For a holomorphic family $\pi : V \rightarrow T$, denote $\mathcal{T}_\pi := (\text{SC}^*)^{-1}(\pi)$. This $\mathcal{T}_\pi$ is regarded as a holomorphic family of isolevel sets.

For any $\varepsilon \in T$, the isolevel set $\mathcal{T}_\pi|_\varepsilon$ can be embedded into $\text{Jac}(\pi^{-1}(\varepsilon)) \times (\mathbb{C}^\times)^{d-1}$ (theorem 2.10). Combining these maps, we obtain the inclusion $\eta^* : \mathcal{T}_\pi \rightarrow \text{Hom}(T, J \times (\mathbb{C}^\times)^{d-1})$ induced from η .

4.1.2 ultradiscretization of η

We are interested in the real positive part of the isolevel set: $\mathcal{V}(\alpha)^+ := \mathcal{V}(\alpha) \cap \text{Hom}(T, \mathbb{R}_+^{N \times M})$. Let $\mathcal{T}_\pi^+ := \mathcal{T}_\pi \cap \text{Hom}(T, \mathbb{R}_+^{N \times M})$. Denote by $\eta_+^* : \mathcal{T}_\pi^+ \rightarrow \text{Hom}(T, J \times (\mathbb{C}^\times)^{d-1})$ the restriction of η^* to $\mathcal{T}_\pi^+$. Let $\mathcal{L} : \text{Hom}(T, J \times (\mathbb{C}^\times)^{d-1}) \rightarrow J(\Gamma) \times \mathbb{R}^{d-1}$ be the mapping defined by

$$((\mathbf{v}(\varepsilon) \bmod (\mathbb{Z}^g + B(\varepsilon)\mathbb{Z}^g), B(\varepsilon)), \mathbf{w}(\varepsilon)) \mapsto (-2\pi i \lim_{\varepsilon \rightarrow 0^+} \varepsilon \mathbf{v}(\varepsilon), - \lim_{\varepsilon \rightarrow 0^+} \varepsilon \log \mathbf{w}(\varepsilon)).$$

By theorem 3.5, this mapping is well-defined.

Our aim in this section is to prove the following theorem:

Theorem 4.1 *There exists a mapping $\text{UD}(\eta)$ such that the following diagram is commutative:*

$$\begin{array}{ccc} \mathcal{T}_\pi^+ & \xrightarrow{\eta_+^*} & \text{Hom}(T, J \times (\mathbb{C}^\times)^{d-1}), \\ \downarrow \text{UD} & & \downarrow \mathcal{L} \\ \mathbb{R}^{N \times M} & \xrightarrow{\text{UD}(\eta)} & J(\Gamma) \times \mathbb{R}^{d-1} \end{array}$$

where Γ is the tropical curve associated with Φ .

First, we refer to the result proved by Mada, Idzumi and Tokihiro [MIT]:

Theorem 4.2 (i) Let $L_n = \text{diag}(V_{n,1}, \dots, V_{n,M}) + S$ and $X := L_N \cdots L_2 L_1$. Then, all the coefficients of $\Phi(x, y) = \det(X(y) - xE)$ are expressed as $\pm(\text{totally positive polynomial in } V_{n,m} \text{ (} n = 1, \dots, N, m = 1, \dots, M))$.
(ii)¹ Coefficients of any (i, j) -minor $\Delta_{i,j}(x, y)$ of $(X(y) - xE)$ are expressed as $\pm(\text{totally positive polynomial in } V_{n,m})$. $\blacksquare$

Proof of theorem 4.1. We prove the theorem by constructing $\text{UD}(\eta)_1 : \mathbb{R}^{N \times M} \rightarrow J(\Gamma)$ and $\text{UD}(\eta)_2 : \mathbb{R}^{N \times M} \rightarrow \mathbb{R}^{d-1}$ respectively.

(I) Constructing UD_1 : For a holomorphic family $\pi : V \rightarrow T$ of analytic curves of genus g , we denote by $\text{Div}^g(\pi) : V_g \rightarrow T$ the holomorphic family of divisors of degree g . A fiber of $\text{Div}^g(\pi)$ is of the form $\{\text{Div}^g(\pi)\}^{-1}(\varepsilon) = \text{Div}^g(\pi^{-1}(\varepsilon))$. By theorem 2.3, we have the mapping $\tilde{\varphi}_C : \mathcal{R}_C = \mathcal{T}_C / \sim \rightarrow \text{Div}^g(C)$ for any spectral curve C . Therefore, this mapping $\tilde{\varphi}_C$ induces a new mapping $\tilde{\varphi}^* : \mathcal{T}_\pi \rightarrow \text{Div}^g(\pi)$.

Let $v = (V_{m,n}(\varepsilon))_{m,n}$ be an element of $\mathcal{T}_\pi$. Denote $X_v = L_N \cdots L_2 L_1$, $L_n = \text{diag}(V_{n,1}, \dots, V_{n,M}) + S$. By theorem 2.3 (ii), x, y coordinates of g points which belongs to $\tilde{\varphi}^*(X_v)(\varepsilon)$ are expressed as roots of polynomials defined by minors of $(X_v - xE)$. Due to theorem 4.2, all the coefficients of these polynomials are of the form $\pm(\text{totally positive polynomials in } V_{n,m})$. Therefore, the limits $\{-\lim_{\varepsilon \rightarrow 0^+} \varepsilon \log x_i(\varepsilon)\}_{i=1}^g$, $\{-\lim_{\varepsilon \rightarrow 0^+} \varepsilon \log y_i(\varepsilon)\}_{i=1}^g$, $(x_i, y_i = \{x, y \text{ coordinates of a point } p_i(\varepsilon) \text{ where } \tilde{\varphi}^*(X_v)(\varepsilon) = p_1 + \cdots + p_g\})$ depend only on $\text{UD}(X(\varepsilon))$. Let P_i be a point in Γ such that $p_i \rightarrow P_i$ (§3.3) and $p_0 : T \rightarrow V$ be an arbitrary section and $p_0 \rightarrow P_0 \in \Gamma$. By theorem 3.6, it is sufficient to define $\text{UD}(\eta)_1(X) := \sum_{i=1}^g F_{P_0}(P_i)$, where $X \in \mathbb{R}^{M \times N}$.

(II) Constructing $\text{UD}(\eta)_2$: We construct $\text{UD}(\eta)_2$ by ultradiscretizing the mapping $i : \mathcal{T}_C \rightarrow (\mathbb{C}^\times)^{d-1}$ introduced in §2.4. By the proof of lemma 2.8, the image $i(X)$ is expressed as $[a_1 : \cdots : a_d]$, where any a_i is a root of some polynomials of which coefficients are components of X . Therefore, the expression $\text{UD}(\eta)_2(X) := (-\lim_{\varepsilon \rightarrow 0^+} \varepsilon \log |a_i|)_i$, $X = \text{UD}(X(\varepsilon))$ is well-defined by example 3.2. $\blacksquare$

By the analogy of the discrete case [MM], we can make a conjecture concerning the ultradiscretized $\tilde{\eta}$.

Conjecture 4.3 The mapping $\text{UD}(\eta)$ is injective.

4.2 fundamental cycles of 2dBBS

Now we consider the 2dBBS (§1). Let $\{W_{n,m}^t, Q_{n,m}^t\}_{n,m}^t$ be a set of real numbers which satisfies the 2dBBS (5). The *fundamental cycle of 2dBBS* is the minimum positive integer F such that

$$\{W_{n,m}^F, Q_{n,m}^F\}_{n,m} = \{W_{n,m}^0, Q_{n,m}^0\}_{n,m}.$$

Let $V_{n,m}^0 = V_{n,m}^0(\varepsilon)$ be a real function satisfying $\text{UD}(V_{n,m}^0) = W_{n,m}^0$. Using the procedure in §2.1, we obtain the matrix function $X(y)(\varepsilon) \in \text{Hom}(T, \text{Mat}(M, \mathbb{C}[y]))$ and therefore the family of spectral curves $\pi : V \rightarrow T$. Denote by Γ the tropical curve defined by $X(y)(\varepsilon)$.

Here we recall proposition 2.6. We denote by $p_0(\varepsilon)$, $p_1(\varepsilon)$, $p_2(\varepsilon)$, $p_3^{(m)}(\varepsilon)$ the points in $C_\varepsilon = \pi^{-1}(\varepsilon)$ as in proposition 2.6

Proposition 4.4 Let $P_0, P_1, P_2, P_3^{(m)} \in \Gamma$ be the tropicalization of $p_0, p_1, p_2, p_3^{(m)}$. ($p_i^{(m)} \rightarrow P_i^{(m)}$). Then, we have $P_0 : (X, Y) = (-\infty, -\infty)$, $P_1 : (X, Y) = (G, A)$, $P_2 : (X, Y) = (+\infty, B)$, $P_3^{(m)} : (X, Y) = (H_m, +\infty)$, where $A = \sum_{m=1}^M Q_{n,m}^0$, $B = \sum_{m=1}^M W_{n,m}^0$, $H_m = \sum_{n=1}^N W_{n,m}^0$ and G is the maximum number such that $(G, A) \in \Gamma$.

¹In the paper [MIT], only the proof of the part (i) is given. However, the part (ii) is also proved by the exactly same argument.

Proof. $\text{UD}(\alpha) = A$, $\text{UD}(\beta) = B$ and $\text{UD}(\gamma_m) = H_m$ are obtained by definition. The G is obtained from $G = \text{UD}(\kappa)$, where κ is the complex number in lemma 2.5 (ii). $\blacksquare$

Using Γ , we can calculate the tropical period matrix B_Γ and vectors $\vec{T} := F_{P_1}(P_0)$, $\vec{N} := F_{P_2}(P_0)$, $\vec{M}^{(m)} := F_{P_3^{(m)}}(P_0) \in J(\Gamma)$ by combinatorial procedures.

Proposition 4.5 *Let F'' be the minimum positive integer such that $F'' \cdot \vec{T} \equiv 0$ and $F' := \text{l.c.m.}(F'', d)$. The fundamental cycle F is a multiple of F' . If conjecture 4.3 is true, then $F = F'$.*

Proof. This proposition is the consequence of theorems 2.10, 3.6, 4.1. $\blacksquare$

Note that the condition $F \cdot \vec{T} \equiv 0 \in J(\Gamma)$ is equivalent to $F \cdot B_\Gamma^{-1} \vec{T} \in \mathbb{Z}^g$.

4.2.1 example I

Let $e := e^{-1/\varepsilon} \in \text{Hom}(T, \mathbb{R}_+)$. Consider the following example:

$$\begin{array}{cccc} 1|..1 & .|1.. & .|.1. & 1|..1 \\ .|.1. & 1|..1 & .|1.. & .|.1. \\ .|1.. & .|.1. & 1|..1 & .|1.. \\ \mathbf{t}=0 & \mathbf{t}=1 & \mathbf{t}=2 & \mathbf{t}=3 \end{array}$$

and $A = B = 1$, $M = N = 3$, $d = 3$. From the picture, we have:

$$X_1^0 = L_3^0 L_2^0 L_1^0 = (\text{diag}(e^1, e^0, e^0) + S)(\text{diag}(e^0, e^1, e^0) + S)(\text{diag}(e^0, e^0, e^1) + S).$$

Its characteristic polynomial $\Phi(x, y)$ satisfies

$$\Phi(x, y) = \det(X_1^0(y) - xE) = -x^3 + x^2(3y + 3e) - x(3y^2 - 21ey + 3e^2) + (y + e)^3.$$

Therefore, the tropical curve Γ defined by Φ satisfies $B_\Gamma = 0$, which implies $J(\Gamma) = \{0\}$. By proposition 4.5, the fundamental cycle is a multiple of $d = 3$. (In fact, the fundamental cycle is 3.)

4.2.2 example II

Start with the following initial value:

$$\begin{array}{ccccc} .|.11 & .|.2. & .|1..1 & 1|...2 & .|.11 \\ .|.11. & .|.2.. & .|.11 & .|.2. & ...|.11. \\ 1|21.1 & 1|2..2 & 1|121. & .|22.. & 1|21.1 \\ \mathbf{t}=0 & \mathbf{t}=1 & \mathbf{t}=2 & \mathbf{t}=3 & \mathbf{t}=8 \end{array}$$

and $A = 1$, $B = 2$, $M = 3$, $N = 4$, $d = 1$. Form the picture, we have:

$$X_1^0 = (\text{diag}(e^2, 1, 1) + S)(\text{diag}(e, e, 1) + S)(\text{diag}(1, e, e) + S)(\text{diag}(e, 1, e) + S).$$

Then $\Phi(x, y) = -x^3 + x^2(5y + 2e^2) - x(y^2 - 7e^2y + e^4) + (y + e^2)^4 + (\text{small})$. Therefore, the tropical curve Γ defined by Φ is as figure 5:

Let $O = (0, 0)$, $v_1 = (2, 2)$ and $v_2 = (4, 2)$. Denote two edges connecting O and v_1 by e_1 and e_2 , and two edges connecting v_1 and v_2 by e_3 and e_4 . Define closed paths $\beta_1, \beta_2, \beta_3$ on Γ as follows:

$$\beta_1 : v_1 \xrightarrow{e_1} O \xrightarrow{e_2} v_1, \quad \beta_2 : v_2 \xrightarrow{e_3} v_1 \xrightarrow{e_4} v_2, \quad \beta_3 : O \rightarrow v_2 \xrightarrow{e_4} v_1 \xrightarrow{e_2} O.$$

Therefore, the tropical period matrix is $B_\Gamma = \begin{pmatrix} 4 & 0 & -2 \\ 0 & 4 & -2 \\ -2 & -2 & 6 \end{pmatrix}$. By proposition 4.4, we have $P_0 =$

$(-\infty, -\infty)$, $P_1 = (2, 1)$, $P_2 = (+\infty, 2)$, $P_3^{(1)} = (4, +\infty)$ and $P_3^{(2)} = P_3^{(3)} = (2, +\infty)$. Then, $\vec{T} =$




Figure 5: Tropical curve with multiplicity $\Gamma \rightarrow \Gamma^0$ in Example II.

$(0, 0, -1)^T$, $\vec{N} = (0, 0, -2)^T$, $\vec{\mathcal{M}}^{(1)} = (2, 2, 0)^T$, $\vec{\mathcal{M}}^{(2)} = \vec{\mathcal{M}}^{(3)} = (2, 0, 0)^T$. It follows that $(B_\Gamma)^{-1}\mathbf{v}_\mathcal{T} = (-1/8, -1/8, -1/4)^T$, which implies that the fundamental cycle is a multiple of 8. (In fact, the fundamental cycle is 8). We note the relations $4\vec{N} = B_\Gamma(-1, -1, -2)^T \in B_\Gamma\mathbb{Z}^3$ and $\mathbf{v}_\mathcal{M}^{(1)} + \mathbf{v}_\mathcal{M}^{(2)} + \mathbf{v}_\mathcal{M}^{(3)} = B_\Gamma(2, 1, 1)^T \in B_\Gamma\mathbb{Z}^3$, which reflect the condition $X_{n+4,m}^t = X_{n,m+3}^t = X_{n,m}^t$.

Acknowledgment

The author is very grateful to Professor Tetsuji Tokihiro and Professor Ralph Wilcox for helpful comments on this paper. Akane Nakamura gave the author a hint for the proof of proposition 2.2. The author appreciates the dedicated support by Kayo Ejiri. This work was supported by KAKENHI 23-1939.

A Proof of lemmas

A.1 Proof of lemma 2.8

Let C be the spectral curve defined in §2.1, p_0 be the unique point contained in $C \setminus C_0$ and k be the local coordinate around p_0 . Let $S_k := (\delta_{i+1,j}) + k^{-M}E_{M,1}$. At p_0 , the equation $X\mathbf{v} = x\mathbf{v}$ implies $k^N(L_N + S_k) \cdots (L_2 + S_k)(L_1 + S_k)\mathbf{v} = (1 + O(k))\mathbf{v}$.

Lemma A.1 (lemma 2.8) *Up to multiplication by a constant, the i -th component f_i of $\mathbf{v}$ has the following expression near p_0 :*

$$f_i = a_i k^{M-i} + o(k^{M-i}), \quad a_i \neq 0.$$

Moreover, we have $a_{i+d} = a_i$ for all i .

Proof. Let $d = \text{g.c.d.}(N, M)$, $N = dN_1$ and $M = dM_1$. Denote $Y_{p,q} := (y_{pd+i, qd+j})_{i,j=1}^d$ for a square matrix $Y = (y_{i,j})_{i,j=1}^M$ of size M . Let $K = (\kappa_{i,j})$ be the diagonal matrix such that $(p-1)d < i \leq pd \Rightarrow \kappa_{i,i} = k^{M-pd}$, $p = 1, 2, \dots, M_1$. Putting $Y := k^N K^{-1}(L_N + S_k) \cdots (L_2 + S_k)(L_1 + S_k)K$ and $\mathbf{w} := K^{-1}\mathbf{v}$, we obtain $Y\mathbf{w} = (1 + c_1 k + o(k))\mathbf{w}$. (c_1 is a non-zero constant). Because the (i, j) -element $x_{i,j}$ of $(L_N + S_k) \cdots (L_2 + S_k)(L_1 + S_k)$ satisfies $x_{i,j} \in \mathbb{C}[k]$ ($i \leq j$), $x_{i,j} \in k^{-M}\mathbb{C}[k]$ ($i > j$) and $x_{i,j+N} = 1$ ($x_{i,j+M} \equiv x_{i,j}k^{-M}$), the (p, q) -th block $Y_{p,q}$ satisfies

$$Y_{p,q} = \begin{cases} E + (\text{strictly lower triangle mat.}) + O(k^d), & q - p \equiv N_1 \pmod{M_1}, \\ O(k^d), & \text{otherwise.} \end{cases}$$

Because $\text{g.c.d.}(N_1, M_1) = 1$, the vector $\mathbf{w}$ must be expressed as:

$$\mathbf{w} = (\overbrace{(0, 0, \dots, 0, 1)}^d, \overbrace{(0, 0, \dots, 0, 1)}^d, \dots, \overbrace{(0, 0, \dots, 0, 1)}^d)^T + O(k^d). \quad (21)$$

On the other hand, for the matrix $Z = Y^{M_1}$, the (p, q) -th block $Z_{p,q}$ satisfies

$$Z_{p,q} = \begin{cases} Y_{p,p+N_1} Y_{p+N_1,p+2N_1} Y_{p+2N_1,p+3N_1} \cdots Y_{p+(M_1-1)N_1,p} & p = q, \\ O(k^d) & p \neq q, \end{cases} \quad (22)$$

where $Y_{p+M_1,q} \equiv Y_{p,q+M_1} \equiv Y_{p,q}$. Then the equation $Y\mathbf{w} = (1 + c_1k + o(k))\mathbf{w}$ induces

$$Z_{p,p}\mathbf{w}_p \equiv (1 + M_1c_1k + o(k))\mathbf{w}_p \pmod{k^d}, \quad p = 1, 2, \dots, M_1, \quad (23)$$

where $\mathbf{w} = (g_i)_{i=1}^M$, $\mathbf{w}_p = (g_{d(p-1)+i})_{i=1}^d$. From this, there exist non-zero complex numbers $a_1^{(p)}, \dots, a_d^{(p)}$ such that

$$\mathbf{w}_p = (a_1^{(p)}k^{d-1}, a_2^{(p)}k^{d-2}, \dots, a_d^{(p)}) + (\text{smaller when } k \rightarrow 0). \quad (24)$$

From (21, 24), we obtain $g_{d(p-1)+i} = a_i^{(p)}k^{d-i} + o(k^{d-i})$ and $a_d^{(1)} = \dots = a_d^{(M_1)}$. Using the equation $\mathbf{v} = K\mathbf{w}$, we rewrite these equations as $f_i = a_i k^{M-i} + o(k^{M-i})$ and $a_d = a_{2d} = \dots = a_{M-d}$.

We prove the equation $a_{i+d} = a_i$ through the following two steps:

Step 1. Let $z_{i,j}^{(p)}$ be the (i, j) -component of the matrix $Z_{p,p}$. We will prove $z_{i+1,i}^{(1)} \equiv z_{i+1,i}^{(2)} \equiv \dots \equiv z_{i+1,i}^{(M_1)} \pmod{k^d}$. Let Y_p^- be a lower triangle matrix with diagonal elements 1 such that $Y_{p,p+N_1} \equiv Y_p^- \pmod{k^d}$. Let $T := (\delta_{i,j+1})_{i,j=1}^d$. Then the matrix Y_p^- is decomposed uniquely as $Y_p^- = E + \sum_{i=1}^{d-1} U_i^{(p)} T^i$, ($U_i^{(p)}$: diagonal). Therefore, we have $Y_{p_1}^- Y_{p_2}^- \cdots Y_{p_l}^- = I + (U_1^{(p_1)} + U_1^{(p_2)} + \dots + U_1^{(p_l)})T + \dots$, which implies that the $(i+1, i)$ -component of the matrix $Y_{p_1}^- Y_{p_2}^- \cdots Y_{p_l}^-$ does not depend on the order of p_i . Because $\{p, p+N_1, \dots, p+(M_1-1)N_1\} \equiv \{1, 2, \dots, M_1\} \pmod{M_1}$ for all p , we conclude the claim of this step by (22).

Step 2. Prove $z_{i+1,i}^{(p)} a_i^{(p)} \equiv M_1 c_1 a_{i+1}^{(p)} \pmod{k^d}$. This is a direct consequence of (23).

From these two steps, we have $a_i^{(p)} = a_i^{(p+1)}$ which is equivalent to $a_i = a_{i+d}$. ■

A.2 Proofs of propositions 2.1, 2.2.

Let $I_n^t = \text{diag}(I_{n,1}^t, \dots, I_{n,M}^t)$, $V_n^t = \text{diag}(V_{n,1}^t, \dots, V_{n,M}^t)$. Then the equation $X_n^t = L_{n+N-1}^t \cdots L_{n+1}^t L_n^t$ is rewritten as

$$X_n^t = Y_0^t + S Y_1^t + \dots + S^{N-1} Y_{N-1}^t + S^N = Z_0^t + Z_1^t S + \dots + Z_{N-1}^t S^{N-1} + S^N,$$

where Y_n^t, Z_n^t ($n = 0, 1, \dots, N-1$) are diagonal matrices. We note that the matrices $Y_0^t, \dots, Y_{N-1}^t$ are independent as functions of $\{V_{n,m}^t\}_{k,m}$.

Proposition A.2 (proposition 2.1) *For given $X_n^t \in \mathcal{W}(\beta)$, there exist M pairs $(R^{(1)}, X^{(1)}), \dots, (R^{(M)}, X^{(M)})$ such that (i) $R^{(i)} = \text{diag}(I_1^{(i)}, \dots, I_M^{(i)}) + S$, (ii) $\alpha = \prod_{m=1}^M I_m^{(i)}$, (iii) $X^{(i)} \in \mathcal{W}(\beta)$, (iv) $X^{(i)} R^{(i)} = R^{(i)} X_n^t$.*

Proof. It is sufficient to prove for a generic X_n^t . In this proof, we denote $I = I_m^{(i)}$, $Y_n = Y_n^t$, $Z_n = Z_n^{t+1}$ for simplicity. Then the equation (8) is rewritten as

$$(Z_0 + Z_1 S + \dots + Z_{N-1} S^{N-1} + S^N)(I + S) = (I + S)(Y_0 + \dots + S^{N-1} Y_{N-1} + S^N),$$

which implies $Z_k S^k I + Z_{k-1} S^k = I S^k Y_k + S^k Y_{k-1}$, ($k = 0, 1, \dots, N$), $Z_{-1} = Y_{-1} = 0$, $Z_N = Y_N = E$. Using these relations recursively and deleting Z_k , we have

$$0 = Y_0 - Z_0 = [Y_0 - S Y_0 S^{-1}] - I(S Y_1 S^{-1}) + (S I S^{-1}) Z_1 = \dots = \sum_{k=0}^N (-1)^k (\Theta_k - S \Theta_k S^{-1}),$$

where $\Theta_k = I(SIS^{-1})(S^2IS^{-2}) \cdots (S^{k-1}IS^{-k+1})(S^kY_kS^{-k})$. This implies that the diagonal matrix $\mathcal{M} := \sum_{k=0}^M (-1)^k \Theta_k$ must satisfy $\mathcal{M} = S\mathcal{M}S^{-1}$, which is equivalent to $\mathcal{M} = \kappa E$ for a certain constant κ . Let $I = -\text{diag}(x_1, \dots, x_M)$ and $S^i Y_i S^{-i} = \text{diag}(c_{i,1}, c_{i,2}, \dots, c_{i,M})$. Then $\mathcal{M} = \kappa E$ is rewritten as

$$\kappa = c_{0,m} + \sum_{i=1}^{N-1} c_{i,m} x_m x_{m+1} \cdots x_{m+i-1} + x_m x_{m+1} \cdots x_{m+N-1} \quad (25)$$

for all $m \in \{1, 2, \dots, M\}$ ($x_{m+M} \equiv x_m$). Let μ_i be a new variable defined by $x_i = \frac{\mu_{i+1}}{\mu_i}$. Then, we have $\frac{\mu_{M+i}}{\mu_i} = x_1 x_2 \cdots x_M = (-1)^M \alpha$ for any i . Therefore, (25) is rewritten as $\kappa = \frac{\sum_{i=0}^{N-1} c_{i,m} \mu_{m+i} + \mu_{m+N}}{\mu_m}$, $\forall m$. Putting $S_\alpha := S|_{y=(-1)^M \alpha}$ and $\boldsymbol{\mu} := (\mu_1, \dots, \mu_M)^T$, we obtain $(Y_0 + S_\alpha Y_1 + S_\alpha^2 Y_2 + \cdots + S_\alpha^{N-1} Y_{N-1} + S_\alpha^N) \boldsymbol{\mu} = \kappa \boldsymbol{\mu}$, which implies

$$(X_n^t|_{y=(-1)^M \alpha}) \boldsymbol{\mu} = \kappa \boldsymbol{\mu}. \quad (26)$$

Because X_n^t is generic, there exist M eigenvectors $\boldsymbol{\mu}$ with non-zero elements. ■

To prove proposition 2.2, we review the well-known result on linear algebra:

Lemma A.3 (the Perron-Frobenius theorem) *Let X be a positive matrix. Then, (i) X has positive eigenvalues, (ii) the maximum eigenvalue $\kappa_{\max}$ in absolute value is positive and simple, (iii) any positive eigenvector of X belongs to $\kappa_{\max}$.* ■

Proposition A.4 (Proposition 2.2) *Let*

$$X_n^t = L_{n+N-1} \cdots L_{n+1} L_n \in \mathcal{W}(\beta), \quad L_n^t = \text{diag}(V_{n,1}^t, \dots, V_{n,M}^t) + S \quad \text{such that} \quad V_{n,m}^t > 0.$$

Then, there uniquely exist two matrices X_n^{t+1} , R_n^t such that (i) $R_n^t = \text{diag}(I_1^t, \dots, I_M^t) + S$, (ii) $\alpha = \prod_{m=1}^M I_m^t$, (iii) $X_n^{t+1} \in \mathcal{W}(\beta)$, (iv) $X_n^{t+1} R_n^t = R_n^t X_n^t$, (v) $I_{n,m}^t > 0$.

Proof. Let us recall the inequality: $\alpha > \beta > 0$. Denote $X_\alpha := X_n^t|_{y=(-1)^M \alpha}$, $S_\alpha := S|_{y=(-1)^M \alpha}$ and $V_n = \text{diag}(V_{n,1}^t, \dots, V_{n,M}^t)$. We start with (26): $\kappa^{-1} \boldsymbol{\mu} = X_\alpha^{-1} \boldsymbol{\mu}$. It is sufficient to prove the existence and the uniqueness of a real eigenvector $\boldsymbol{\mu} = (\mu_1, \dots, \mu_M)^T$ such that $(-1)^{k+1} \mu_k > 0$ for all k . (Note that $I_{n,i}^t = -\mu_{i+1}/\mu_i$). The vector $\boldsymbol{\mu}$ must satisfy $P \boldsymbol{\mu} > 0$, where $P = \text{diag}(1, -1, 1, \dots, (-1)^M)$. By (7), we have

$$X_\alpha^{-1} = (E + S_\alpha^{-1} V_n)^{-1} S_\alpha^{-1} (E + S_\alpha^{-1} V_{n+1})^{-1} S_\alpha^{-1} \cdots (E + S_\alpha^{-1} V_{n+N-1})^{-1} S_\alpha^{-1}.$$

By the inequality $\alpha > \beta > 0$, $V_{n,m}^t > 0$, we have the following expression:

$$(E + S_\alpha^{-1} V_{n'})^{-1} = E - S_\alpha^{-1} V_{n'} + (S_\alpha^{-1} V_{n'})^2 - (S_\alpha^{-1} V_{n'})^3 + (S_\alpha^{-1} V_{n'})^4 - \cdots,$$

which implies that the (i, j) -element of $(E + S_\alpha^{-1} V_{n'})^{-1}$ is positive (*resp.* negative) if $i + j$ is even (*resp.* odd). In other words, $P(E + S_\alpha^{-1} V_{n'})^{-1} P$ is a positive matrix. Therefore $(-1)^N P X_\alpha^{-1} P$ is also a positive matrix. The proposition follows from this fact and the Perron-Frobenius theorem. ■

A.3 Proof of lemma 2.5

In this section, we calculate the determinants of matrices introduced in §2.3. First, the equations $\det R_n^t = \alpha - (-1)^M y$, $\det L_n^t = \beta - (-1)^M y$ are straightforward.

Next, we recall the equation (16). The $z_{i,j}$ in (16) should satisfy

$$z_{i,j} + y z_{i,j+M} + y^2 z_{i,j+2M} + \cdots = \{(i, j)\text{-th element of } X_{n,m}^t(y)\}.$$

Define two vectors $\mathbf{I}$ and $\boldsymbol{\mu}$ by

$$\mathbf{I} = (1, -I_{n,1}^t, I_{n,1}^t I_{n,2}^t, \dots, \prod_{m=1}^{N-1} (-I_{n,m}^t))^T, \quad \boldsymbol{\mu} = (1, -I_{n,1}^t, I_{n,1}^t I_{n,2}^t, \dots, \prod_{m=1}^{M-1} (-I_{n,m}^t))^T.$$

Then, we have

$$(z_{1,1}, z_{1,2}, \dots, z_{1,N}) \cdot \mathbf{I} = (1, 0, 0, \dots, 0) \cdot (X_{n,m}^t|_{y=(-1)^{M\alpha}}) \cdot \boldsymbol{\mu}.$$

By (26), this equals to κ . We hence have $M_{n,m}^t \mathbf{I} = (0, \dots, 0, x - \kappa)^T$. On the other hand, due to the definition of M_n^t , $\det M_n^t$ must be a monic or anti-monic polynomial in x of degree 1. Therefore, we have $\det M_n^t = (-1)^{N+1}(x - \kappa)$.

We calculate $\det H_n^t$ similarly. Let

$$\mathbf{V} = (1, -V_{n,1}^t, V_{n,1}^t V_{n,2}^t, \dots, \prod_{m=1}^{N-1} (-V_{n,m}^t))^T, \quad \boldsymbol{\nu} = (1, -V_{n,1}^t, V_{n,1}^t V_{n,2}^t, \dots, \prod_{m=1}^{M-1} (-V_{n,m}^t))^T.$$

Then, we have

$$(z_{1,1}, z_{1,2}, \dots, z_{1,N}) \cdot \mathbf{V} = (1, 0, 0, \dots, 0) \cdot (X_{n,m}^t|_{y=(-1)^{M\beta}}) \cdot \boldsymbol{\nu} = 0,$$

which implies $H_{n,m}^t \mathbf{V} = (0, \dots, 0, x)^T$. Therefore, $\det H_{n,m}^t = (-1)^{N+1}x$.

At last, we can directly calculate $\det U_m^t = (-1)^M(x - z_{1,1}) = (-1)^{N+1}(x - V_{N,m}^t \cdots V_{2,m}^t V_{1,m}^t)$. ■

References

- [HHIKTT] Hatayama G, Hikami K, Inoue R, Kuniba A, Takagi T and Tokihiro T, *J. Math. Phys.* **42** (2001) 274
- [HIK] Hikami K, Inoue R and Komori Y, *J. Phys. Soc. Jpn.* **68** (1999) 2234-2240
- [IH] Inoue R and Hikami K, *J. Phys. A: Math. Gen.* **32** (1999) 6853
- [IMS] Itenberg I, Mikhalkin G and Shustin E I, “Tropical algebraic geometry”, *Overwolfach Seminars* Birkhauser (Architectural), Berlin (2007)
- [I1] Iwao S, *Int. Math. Res. Notices* **2010** (2010) 112–148
- [I2] Iwao S, *J. Phys. A: Math. Theor.* **43** (2010) 155208
- [KNY] Kajiwara K, Noumi M and Yamada Y, *Lett. Math. Phys.* **60** (2002) no. 3, 211–219
- [MIT] Mada J, Idzumi M and Tokihiro T, *J. Math. Phys.* **46** (2005) 022701
- [M] Mikhalkin G, Tropical geometry and its applications, *Preprint* math/06011041
- [MZ] Mikhalkin G and Zharkov I, Tropical curves, their jacobians and theta functions, *Preprint* math/0612267
- [MM] Morebeke P and Mumford D, *Acta. Math.* **143** (1978) 93–154
- [MNSTTTM] Moriwaki S, Nagai A, Satsuma J, Tokihiro T, Torii M, Takahashi D and Matsukidaira J, *Symmetries and Integrability of Difference Equations*, London Math. Soc. Lecture Note Series 255 (ed: Peter A. Clarkson and Frank W. Nijhoff) Cambridge (1999) 334–342
- [TS] Takahashi D and Satsuma J, *J. Phys. Soc. Jpn.* **59** (1990) 3514–3519
- [TM] Takahashi D and Matsukidaira J, *J. Phys. A: Math. Gen.* **30** (1997) L733
- [TTMS] Tokihiro T, Takahashi D, Matsukidaira J and Satsuma J, *Phys. Rev. Lett.* **76** (1996) 3247–3250
- [TNS] Tokihiro T, Nagai A and Satsuma J, *Inverse Problems* **15** (1999) 1639